

Response of the Corpus Luteum to Luteinizing Hormone

by G. D. Niswender*

The response of steroidogenic tissues to tropic hormones is regulated in part by specific receptors in the target cells for the stimulatory hormone. As a result of hormone binding to receptor the enzyme adenylate cyclase is activated with a resultant increase in intracellular levels of cAMP. Enhanced protein kinase activity then leads to increased steroidogenesis via several possible mechanisms, including direct activation of components of steroidogenic enzyme systems via phosphorylation.

The initial effects of tropic hormones such as LH are dependent upon the number of receptors present on the surface of the target cell. Numerous factors influence the number of LH receptors in the corpus luteum. A model is presented for the mechanisms involved in the loss and renewal of LH receptors in the luteal cell. The life of the LH receptor on luteal cells appears to be a single binding of hormone. The hormone-receptor complex is then internalized by endocytosis and the hormone is degraded in lysosomes. After internalization the receptor is also degraded in lysosomes or recycled via the Golgi apparatus. New or recycled receptors for LH are incorporated into the limiting membrane of protein containing secretory granules. One of the actions of LH is enhancement of the exocytosis of these secretory granules with incorporation of the limiting membrane (and the LH receptors?) of the granule into the plasma membrane of the cell. These proposed mechanisms explain the increase in the number of receptors for LH seen immediately after stimulation of the luteal cell with massive doses of LH and also explain the "down-regulation" of LH receptors 24 hr after administration of LH.

The purpose of this communication is to describe some of the mechanisms involved in the regulation of ovarian progesterone secretion by luteinizing hormone (LH). Particular attention will be given to the function of receptors for LH and the mechanisms involved in the regulation of LH receptor numbers. Although this discussion will, in general, be limited to the ovarian luteal cell, it is known that most of the mechanisms to be discussed are also operable in the ovarian follicle, the testes and the adrenal gland. Therefore, the concepts to be presented appear to be general and are likely to be applicable to all steroid secreting glands.

In mammals, LH secreted by the anterior pituitary gland appears to be the primary factor which regulates the secretion of progesterone from the corpus luteum (1-5). Other endocrine factors are also important for regulation of the lifespan and

function of the corpus luteum including prolactin in rodents, prostaglandins in domestic animals and rodents and placental or embryonic hormones in most species.

The actions of LH at the level of the luteal cell appear to be mediated via a specific receptor for this hormone in the plasma membrane (5-7). Interaction of LH with its specific receptors results in activation of adenylate cyclase and results in elevated intracellular levels of cAMP (8). The elevated cAMP results in enhanced activity of protein kinase and increased steroidogenesis (7, 9). There is evidence that the activity of at least two of the enzymes in the steroidogenic pathway in luteal cells is influenced directly by protein kinase. The activities of cholesterol esterase, the enzyme responsible for cleavage of the ester form of cholesterol to allow its utilization as substrate for steroidogenesis (10), and of cholesterol side-chain cleavage complex (11), which is thought to be the rate-limiting enzyme complex in steroid-secreting cells, both appear to be increased as a result of increased protein kinase

*Department of Physiology and Biophysics, Colorado State University, Fort Collins, Colorado 80523.

activity. The effects of LH on steroidogenesis have been reviewed recently (5, 8).

The effects of LH are mediated via a specific receptor in the plasma membrane of target cells. Therefore, the responsiveness of the target tissue is dependent upon the number of receptors present on the cell. Thus, the factors which regulate the numbers of these receptors become important in regulating target cell function. There are several endocrine mechanisms which influence the number of receptors for LH on the luteal cell. First, dramatic elevations of LH itself can have both a positive (up-regulation) and a negative (down-regulation) effect on the number of receptors in target tissues for LH. Following administration of 1 mg of ovine LH to sheep there is a significant increase in the number of receptors for LH within 10 min but by 12 to 24 hr the total number of receptors is decreased to 20-30% of that seen prior to the injection (12). Within 48 hr the number of receptors for LH had returned to preinjection levels. There was a high degree of correlation between the number of receptors occupied by LH at 10 min and the number lost by 24 hr. This observation suggests that loss of receptors for LH is coupled to binding of LH to the receptor.

The phenomenon of "down-regulation" of receptors for LH has also been demonstrated in rat luteal (13), follicular (14), and testicular (15) tissues. This phenomenon is more pronounced in studies using human chorionic gonadotropin (hCG) than when LH is employed due to the much longer life of hCG in the circulation (12). The biological significance of "down-regulation" of receptors for LH is unclear at the present time. In general, it appears to be a "pharmacologic" effect due to administration of massive quantities of exogenous hormone. The single exception may be the loss of LH receptors in follicular granulosa cells during luteinization, after the naturally occurring preovulatory LH surge (16, 17). However, this point is controversial, since Nimrod et al. (18) were unable to detect any significant loss of receptors for LH under physiological conditions.

It is important to point out that the use of "pharmacologic" levels of LH or hCG to study the kinetics of loss and renewal of LH receptors is necessary in order to magnify the relatively minor changes in receptor numbers which occur with the levels of LH which circulate normally. The use of these and other procedures has allowed us to formulate a hypothetical model for the mechanisms which modulate the turnover of LH receptors in the luteal cell membrane (5). It appears that LH binds to its specific receptor which results in activation of the intracellular events leading to enhanced pro-

gesterone secretion (5, 8). The LH-receptor complex is then internalized via endocytosis (19-22) and the hormone is subsequently degraded in lysosomes (23, 24). The receptor is either degraded in the lysosome or recycled. New receptor, synthesized in the endoplasmic reticulum, or recycled receptor is then incorporated into the limiting membrane of protein containing secretory granules at the level of the Golgi apparatus. Under the stimulatory action of LH, or other secretagogues such as beta-agonists or prostaglandin E₂, the release of these granules is stimulated via exocytosis with the limiting membrane of the secretory granule along with the receptors for LH being incorporated into the plasma membrane of the cell (5, 24). This model would predict that stimulation of the luteal cell with large quantities of LH would result in exocytosis of large numbers of secretory granules and increased numbers of LH receptors immediately following stimulation (i.e., 10-30 min). However, after granule numbers are depleted, synthesis of new granules would be required before the process could continue. During this period of granule synthesis, LH-receptor complexes would continue to be internalized resulting in a net loss of receptor (down-regulation) from the plasma membrane. Thus, what is predicted and the experimental data are in good agreement. There is also good evidence that exocytosis of granules is associated with the secretion of progesterone (25).

Other hormones also play a role in the regulation of ovarian receptors for LH. In rats, prolactin appears necessary for normal numbers of receptors for LH in the corpus luteum (26). This role is consistent with the known biological effects of prolactin on luteal function in this species. However, prolactin does not appear to influence luteal function in cattle (3, 27) or sheep (28, 29). Although receptors for prolactin have been demonstrated in porcine luteal tissue (30), the role of this hormone in regulation of luteal function in this species has not been defined clearly. There is also good evidence that estradiol-17 β and follicle-stimulating hormone act synergistically to induce LH receptors in the maturing rat follicular granulosa cell (17).

Finally, there is considerable information which correlates the number of receptors for LH and biological function of the corpus luteum. In rats, receptors for LH appear first in granulosa cells of the maturing follicle (17, 31) and as the follicle matures and increases in size the number of receptors for LH increases dramatically (16, 32-34). Then, after the preovulatory surge of LH the number of granulosa cell receptors for LH decreases. A similar pattern of receptor development has been reported in sheep (35) and pigs (36).

After ovulation the number of luteal receptors for LH appears to decrease (16, 37), although this point is controversial (18). Once the corpus luteum begins to develop, secretion of progesterone is correlated highly with the number of receptors for LH in rats (38, 39), cows (40), sheep (41), and women (42). Since basal levels of LH do not change throughout the luteal phase of the reproductive cycle and since the average association constant of the LH receptor for LH does not change (41) the increased number of total receptors for LH results in an increase in the number of receptors occupied by endogenous hormone. Therefore, there is also an excellent correlation between the number of LH receptors occupied by endogenous hormone and serum levels of progesterone. This was to be expected based on our current understanding of the actions of LH.

As the corpus luteum regresses during the late luteal phase of the reproductive cycle, there is a dramatic decline in the number of receptors for LH and serum levels of progesterone (41). The loss of receptors for LH during luteolysis in rats appears to be mediated by prostaglandin $F_2\alpha$ ($PGF_2\alpha$) (43). However, administration of a luteolytic dose of $PGF_2\alpha$ to ewes during the mid-luteal phase of the estrous cycle resulted in decreased levels of progesterone in serum several hours before a decline in the number of receptors for LH (44). Thus, it appeared that the reduced number of receptors for LH was a result of luteolysis and not the mechanism whereby $PGF_2\alpha$ induced luteal regression.

This brief review has provided an overview of the factors involved in the regulation of LH receptors and progesterone secretion in the luteal cell. For an in-depth examination of the various aspects of these subjects three recent reviews are recommended (5, 7, 8).

REFERENCES

- Short, R. V. Ovarian steroid synthesis *in vivo*. *Rec. Prog. Horm. Res.* 20: 303 (1964).
- Niswender, G. D., Menon, K. M. J., and Jaffe, R. B. Regulation of the corpus luteum during the menstrual cycle and early pregnancy. *Fertil. Steril.* 23: 432 (1972).
- Hansel, W., Concannon, P. W., and Lukaszewska, J. H. Corpora lutea of the large domestic animals. *Biol. Reprod.* 8: 22 (1973).
- Nalbandov, A. V. Control of luteal function in mammals. In: *Handbook of Physiology. Endocrinology II, Part I*. Waverly Press, Baltimore, 1973.
- Niswender, G. D., Sawyer, H. R., Chen, T. T., and Endres, D. B. Action of luteinizing hormone at the luteal cell level. In: *Advances in Sex Hormone Research*, Vol. 4, J. A. Thomas and R. L. Singhal, (Eds.), Urban & Schwarzenberg, Baltimore, 1980.
- Ryan, R. J., and Lee, C. J. The role of membrane bound receptors. *Biol. Reprod.* 14: 16 (1976).
- Dufau, M. L., and Catt, K. J. Gonadotropin receptors and regulation of steroidogenesis in the testis and ovary. *Vit. Horm.* 36: 462 (1978).
- Marsh, J. The role of cyclic AMP in gonadal function. In: *Advances in Cyclic Nucleotide Research*. P. Greengard and G. A. Robison, (Eds.), Raven Press, New York, 1975.
- Ling, W. Y. and Marsh, J. M. Reevaluation of the role of cyclic adenosine 3', 5'-monophosphate and protein kinase in stimulation of steroidogenesis by luteinizing hormone in bovine corpus luteum slices. *Endocrinology* 100: 1571 (1977).
- Caffrey, J. L., Fletcher, P. W., Diekmann, M. A., O'Callaghan, P. L., and Niswender, G. D. The activity of ovine luteal cholesterol esterase during several experimental conditions. *Biol. Reprod.* 21: 601 (1978).
- Caron, M. G., Goldstein, S., Savard, K., and Marsh, J. M. Protein kinase stimulation of a reconstituted cholesterol side chain cleavage enzyme system in the bovine corpus luteum. *J. Biol. Chem.* 250: 5137 (1975).
- Suter, D. E., Fletcher, P. W., Sluss, P. M., Reichert, L. E., Jr., and Niswender, G. D. Alterations in the number of ovine luteal receptors for LH and progesterone secretion induced by homologous hormone. *Biol. Reprod.* 22: 205 (1980).
- Conti, M., Harwood, J. P., Hsueh, A. J. W., Dufau, M. L., and Catt, K. J. Gonadotropin-induced loss of hormone receptors and desensitization of adenylate cyclase in the ovary. *J. Biol. Chem.* 251: 7729 (1977).
- Ryan, R. J., Birnbaumer, L., Lee, C. Y., and Hunzicker-Dunn, M. Gonadotropin interactions with the gonad as assessed by receptor binding and adenylyl cyclase activity. *Int. Rev. Physiol.* 13: 85 (1977).
- Hsueh, A. J. W., Dufau, M. L., and Catt, K. J. Gonadotropin-induced regulation of luteinizing hormone receptors and desensitization of testicular 3': 5'-cyclic AMP and testosterone response. *Proc. Natl. Acad. Sci. (U.S.A.)* 74: 592 (1977).
- Richards, J. S., Ireland, J. J., Rao, M. C., Bernath, G. A., Midgley, A. R., Jr., and Reichert, L. E., Jr. Ovarian follicular development in the rat: Hormone receptor regulation by estradiol, follicle-stimulating hormone and luteinizing hormone. *Endocrinology* 99: 1562 (1976).
- Richards, J. S. Hormone regulation of hormone receptors in ovarian follicular development. In: *Ovarian Follicular Development*. A. R. Midgley and W. A. Sadler, Eds., Raven Press, New York, 1979.
- Nimrod, A., Bedrak, E., and Lamprecht, S. A. Appearance of LH-receptors and LH-stimulable cyclic AMP accumulation in granulosa cells during follicular maturation in the rat ovary. *Biochem. Biophys. Res. Commun.* 78: 977 (1977).
- Chen, T. T., Abel, J. H., Jr., McClellan, M. C., Sawyer, H. R., and Niswender, G. D. Localization of gonadotropic hormones in lysosomes of ovine luteal cells. *Cytobiology* 14: 412 (1977).
- Conn, P. M., Conti, M., Harwood, J. P., Dufau, M. L., and Catt, K. J. Internalization of gonadotropin-receptor complex in ovarian luteal cells. *Nature* 274: 598 (1978).
- Anderson, W., Kang, Y.-H., Perotti, M. E., Bramley, T. A., and Ryan, R. J. Interactions of gonadotropins with corpus luteum membranes. III. Electron microscopic localization of [125 I]-hCG binding to sensitive and desensitized ovaries seven days after PMSG-hCG. *Biol. Reprod.* 20: 362 (1978).
- Amsterdam, A., Kohen, F., Nimrod, A., and Lindner, H. R. Lateral mobility and internalization of hormone receptors to human chorionic gonadotropin in cultured rat granulosa cells. *Adv. Exptl. Med. Biol.* 112: 69 (1979).

23. Ascoli, M., and Puett, D. Intracellular uptake and catabolism of lutropin by testicular tissue *in vivo*. FEBS Letters 75: 77 (1977).
24. Niswender, G. D. Factors regulating LH receptors on ovine luteal cells. J. Reprod. Fertil. Suppl., in press.
25. Sawyer, H. R., Abel, J. H., Jr., McClellan, M. C., Schmitz, M., and Niswender, G. D. Secretory granules and progesterone secretion by ovine corpora lutea *in vitro*. Endocrinology 104: 476 (1979).
26. Richards, J. S., and Williams, J. J. Luteal cell receptor content for prolactin (PRL) and luteinizing hormone (LH): Regulation by LH and PRL. Endocrinology 99: 1571 (1976).
27. Hoffman, B., Schams, D., Bopp, R., Ender, M. L., Gimenez, T., and Karg, H. Luteotrophic factors in the cow: Evidence for LH rather than prolactin. Reprod. Fertil. 40: 77 (1974).
28. Karsch, F. J., Cook, B., Ellicott, A. R., Foster, D. L., Jackson, G. L., and Nalbandov, A. V. Failure of infused prolactin to prolong the life span of the corpus luteum of the ewe. Endocrinology 89: 272 (1971).
29. Niswender, G. D. Influence of 2-Br- α -ergocryptine on serum levels of prolactin and the estrous cycle of sheep. Endocrinology 94: 912 (1974).
30. Rolland, R., Hammond, J. M., Schellekens, L. A., Lequin, R. M., and deJong, F. H. Prolactin and ovarian function. In: The Endocrine Function of the Human Ovary. V.H.T. James, M. Serio, and G. Giusti, Eds., Academic Press, New York, 1976.
31. Channing, C., and Kammerman, S. Binding of gonadotropins to ovarian cells. Biol. Reprod. 10: 179 (1974).
32. Zeleznik, A. J., Midgely, A. R., Jr., and Reichert, L. E., Jr. Granulosa cell maturation in the rat: Increased binding of human chorionic gonadotropin following treatment with follicle stimulating hormone *in vivo*. Endocrinology 95: 818 (1974).
33. Lee, C. Y. The procine ovarian follicle: III. Development of chorionic gonadotropin receptors associated with increase in adenylate cyclase activity during follicle maturation. Endocrinology 99: 42 (1976).
34. Chang, S. C. S., Anderson, W., Lewis, J. C., Ryan, R. J., and Kong Y.-H. The procine ovarian follicle: II. Electron microscopic study of surface features of granulosa cells at different stages of development. Biol. Reprod. 16: 349 (1977).
35. Carson, R. S., Findlay, J. K. and Burger, H. G. Receptors for gonadotrophins in the ovine follicle during growth and atresia. In: Ovarian Follicular and Corpus Luteum Function, Advances in Experimental Medicine and Biology, Vol. 112, C. P. Channing, J. Marsh, and W. A. Sadler, Eds., Plenum Press, New York, 1979.
36. Kammerman, S., and Ross, J. Increase in number of gonadotropin receptors on granulosa cells during follicle maturation. J. Clin. Endocrinol. Metab. 41: 546 (1975).
37. Rao, M. C., Richards, J. S., Midgley, A. R., Jr., and Reichert, L. E. Regulation of gonadotrophin receptors by LH in granulosa cells. Endocrinology 101: 512 (1977).
38. Hacik, T., and Kolena, J. Production of oestradiol, cAMP and ¹²⁵I-hCG binding by rat ovaries during the reproductive cycle. Endocrinol. 66: 15 (1975).
39. Lee, C. Y., Tateishi, K., Ryan, R. J., and Jiang, N. S. Binding of human chorionic gonadotropin by rat ovarian slices: Dependence on the functional state of the ovary. Proc. Soc. Exptl. Biol. Med. 148: 505 (1975).
40. Rao, C. V., Estergreen, V. L., Carman, F. R., Moss, G. E. and Frandle, K. A. Receptors for prostaglandin (PG)F₂ α and human choriogonadotropin (hCG) in cell membranes of bovine corpora lutea (CL) throughout the estrous cycle. V. Int. Cong Endocrin., abstract (1976).
41. Diekman, M. A., O'Callaghan, P. L., Nett, T. M. and Niswender, G. D. Validation of methods and quantification of luteal receptors for LH throughout the estrous cycle and early pregnancy in ewes. Biol. Reprod. 19: 999 (1978).
42. Lee, C. Y., Coulan, C. B., Jiang, N. S., and Ryan, R. J. Receptors for human luteinizing hormone in human corpora luteal tissue. J. Clin. Endocrinol. Metab. 36: 148 (1973).
43. Grinwich, D. L., Hichens, M., and Behrman, H. R. Control of the LH receptor by prolactin and prostaglandin F₂ in rat corpora lutea. Biol. Reprod. 14: 212 (1976).
44. Diekman, M. A., O'Callaghan, P. L., Nett, T. M. and Niswender, G. D. Effect of prostaglandin F₂ α on the number of LH receptors in ovine corpora lutea. Biol. Reprod. 19: 1091 (1978).